

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Annals of Prosthodontics and Restorative Dentistry

Journal homepage: <https://www.aprd.in/>

## Original Research Article

# Assessment of oral, dental and facial pain in patients

Isha Rastogi<sup>1,\*</sup><sup>1</sup>Dept. of Dental, Mayo Institute of Medical Sciences, Barabanki, Uttar Pradesh, India

### ARTICLE INFO

#### Article history:

Received 16-08-2022

Accepted 04-11-2022

Available online 24-11-2022

#### Keywords:

Pain

Dental

Oral

### ABSTRACT

**Introduction:** Discomfort or pain in mouth is a common problem. It needs to be fully catered to and treated at the earliest.

**Aim and Objectives:** To find out causes of oral pain in patients.

**Material and Methods:** 500 patients above 20 years were chosen.

**Results:** Patients have various reasons of oral pain, but they are careless.

**Discussion:** Patients suffer oral pain and some accept it as part of their living.

**Conclusion:** Oral pain should be thoroughly evaluated and taken care of in diagnosis.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Pain in the oral cavity is a serious issue. The International Association for the study of pain defined it as an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage.<sup>1</sup>

Theories of pain perception:<sup>2</sup> are Specificity theory, Pattern theory, Gate control theory. It is exactly pain which is the most common reason for patients to come to the dental clinic; this pain usually originates in the tooth itself or its supporting structures.<sup>3</sup>

Orofacial pain is pain within the trigeminal system.<sup>4</sup> Patients will sometimes present to medical practitioners for the management of pain or other dental or oral problems.<sup>5</sup>

Aim is to inquire patients about dental pain (like conversation and basic tools) and to evaluate oral pain. Objective is to find out about oral pain in rural and urban populations. Clinical relevance is that patients experience oral pain but they do not give much attention to it.

## 2. Materials and Methods

500 subjects above 20 years from village and half from city in 2 months visiting a dental hospital on outer of Malihabad were studied. Personal details were asked and recorded like in normal conversation and then results followed.

## 3. Results

### 3.1. Oral pain

**Table 1:** Descriptive analysis of oral pain in study population (N=500)

Oral pain	N	%
Dental pain	300	60%
Non dental pain	200	40%
Total	500	100%

In this study, it was seen that 60% had dental pain, 40% had non-dental pain. Dental pain comprised of 66.7% that had pulpal pain and 33.3% was of periapical abscess. Non-dental pain was 37.5% of tmj temporomandibular disorders, 15% of sinus, 10% of bruxism, 6% of neurogenic, 4% of

\* Corresponding author.

E-mail address: [excellent123@gmail.com](mailto:excellent123@gmail.com) (I. Rastogi).

**Table 2:** Descriptive analysis (N=500)

<b>Oral pain</b>		<b>N (%)</b>
Dental pain (N=300)	Pulpal pain (Dentinal pain, reversible pulpitis, irreversible pulpitis)	200 (66.7%)
	Periapical abscess	100 (33.3%)
	TMD	75 (37.5%)
	Sinus	30 (15%)
	Bruxism	20 (10%)
Non dental pain (n=200)	Neurogenic	12 (6%)
	Psychogenic	8 (4%)
	Angina referred	5 (2.5%)
	Periapical abscess	50 (25%)

**Table 3:** Descriptive analysis

<b>Age in Years</b>		<b>N (%)</b>
Till 45 Years (N=425)	Pulpal pain (Dentinal pain, reversible pulpitis, irreversible pulpitis)	200 (47.1%)
	Periapical abscess	100 (23.5%)
	TMD	75 (17.6%)
	Bruxism	20 (4.7%)
	Sinus	30 (7.1%)
Above 45 Years (N=75)	Neurogenic	12 (16%)
	Psychogenic	8 (10.7%)
	Angina referred	5 (6.7%)
	Periapical abscess	50 (66.7%)

**Table 4:** Descriptive analysis (N=500)

<b>Gender</b>		<b>N (%)</b>
Male (N=258)	Pulpal pain	94 (36.4%)
	Periapical abscess	52 (20.2%)
	TMD	42 (16.3%)
	Bruxism	11 (4.3%)
	Sinus	18 (7%)
	Neurogenic	7 (2.7%)
	Psychogenic	3 (1.2%)
	Angina referred	53 (20.5%)
	Periapical abscess	28 (10.9%)
	Pulpal pain	116 (47.93%)
Female (N=242)	Periapical abscess	48 (19.8%)
	TMD	33 (13.6%)
	Bruxism	9 (3.7%)
	Sinus	12 (5%)
	Neurogenic	5 (2.1%)
	Psychogenic	5 (2.1%)
	Angina referred	2 (0.8%)
	Periapical abscess	22 (9.09%)

psychogenic, 2.5% of angina referred, 25% of periapical abscess. Till 45 years, 47.1% had pulpal pain, 23.5% had periapical abscess, 17.6% had tmd temporomandibular diseases, 4.7% had bruxism, 7.1% had sinus. Above 45 years, 16% had neurogenic, 10.7% had psychogenic, 6.7% had angina referred and 66.7% had periapical abscess. In males, 36.4% had pulpal pain, 20.2% had periapical abscess, 16.3% had tmd temporomandibular disorders, 4.3% had bruxism, 7% had sinus, 2.7% had neurogenic, 1.2% had psychogenic, 20.5% had angina referred, 10.9% had periapical abscess. In females, 47.93% had pulpal pain, 19.8% had periapical abscess, 13.6% had tmd temporomandibular disorders, 3.7% had bruxism, 5% had sinus, 2.1% had neurogenic, 2.1% had psychogenic, 0.8% had angina referred and 9.09% had periapical abscess.

#### 4. Discussion

Orofacial pain interferes with daily life activities impacting negatively on quality of life.<sup>6</sup> When a patient visits our clinic complaining of dental pain, we diagnose the cause by looking- both visually and radiographically-for organic or functional abnormalities. Nearly all pain is caused by an organic problem such as dental caries, periodontitis, pulpitis or trauma.<sup>7</sup> Dental pain occurs as a result of inflammation of the pulp(pulpitis). This is generally caused by bacteria from decayed teeth or defective dental fillings.<sup>8</sup> Peoples' perceptions of their oral health status and the related impacts of dental pain on their daily lives are important in planning services designed to improve the quality of life of individuals(9).

The orofacial pain classification as outlined by Okeson<sup>9,10</sup> is divided into physical (Axis 1) and psychological (Axis 2) conditions. Physical conditions comprise temporomandibular disorders (TMD), which include disorders of the temporomandibular joint (TMJ) and disorders of the musculoskeletal structures (eg masticatory muscles and cervical spine); neuropathic pains, which include episodic (eg. Trigeminal neuralgia TN) and continuous (eg. Peripheral/centralized mediated) pains and neurovascular disorders (eg. migraine). Psychological conditions include mood and anxiety disorders. Research to date has shown that pain influences most dimensions of quality of life, mainly the physical and emotional ones. QOL is defined as the person's evaluation of his or her well-being and functioning in different life domains.<sup>11</sup>

Questions to ask when assessing oral pain:<sup>12</sup>

When obtaining a pain history, the mnemonic SOCRATES can be useful

1. Site- where is the pain?
2. Onset- when did it start?
3. Character- can you describe the pain?
4. Radiation- does the pain spread anywhere?

5. Associations- are there other problems associated with the pain?
6. Time course- does the pain follow any pattern? How long does it last?
7. Exacerbating or relieving factors- does anything worsen or improve it?
8. Severity- how bad is the pain?

#### 4.1. Assessment of pain<sup>13</sup>

Rating scale techniques are often used. The most commonly used techniques are:

Numerical rating scale, visual analogue scale, McGill pain questionnaire, behavioral rating scale.

The orofacial pain from dental origin was specifically called ‘odontogenic toothache’.<sup>14</sup> Orofacial pain is tenderness in the head, face (including oral cavity) and neck.<sup>15</sup> Pain in the face and mouth is a frequent problem.<sup>16</sup> Facial or orofacial pain refers to any type of pain in the area bounded by the eyes and the lower mandibles, including the oral cavity.<sup>17</sup>

For Patients affected by tmjs- Abdelnabi et al showed that new dentures with corrected occlusion significantly improved clinical signs and symptoms of tmd in complete denture wearers and disc position.<sup>18</sup> Prosthodontic treatment in patients affected by bruxism- various treatments have been suggested based on behaviour modification, such as habit awareness, habit reversal therapy, and relaxation techniques, which may eliminate awake bruxism.<sup>19</sup> The prosthodontist has always played a major role in tmd treatment by providing many different treatments mostly oriented toward prosthetic reconstruction.<sup>20</sup> The instability of the bite and the severity of the clinical presentation need to be considered as possible contraindications for any restorative procedure.<sup>21</sup> Furthermore, we should notice the importance of stable occlusion in the intercuspal position.<sup>22</sup> Management of Orofacial pain can only be effective if the correct diagnosis is reached and may involve referral to secondary or tertiary care.<sup>23</sup> Dental pain is extremely common and it can also coexist with other conditions.<sup>24</sup> Newer disciplines such as oral diagnosis/oral medicine, dental anesthesiology and temporomandibular disorders TMD/Orofacial pain are focused on the diagnosis and treatment of diseases affecting the entire head and neck.<sup>25</sup> The relationship between tmjs and bruxism is controversial.<sup>26</sup> OFP is highly prevalent in the population.<sup>28</sup> Facial pain has a considerable impact on QoL. Different Orofacial pains may cause variable levels of anxiety and depression and various coping strategies, daily limitations or perception of the disease.<sup>27</sup> Moreover recurrent headache has been identified as a neurological disorder also of high prevalence in the general population.<sup>28</sup> Wearing complete dentures does not predispose edentulous individuals to tmjs.<sup>27</sup>

Management of chronic pain conditions is among the most difficult problems confronting clinicians. These

conditions often found in the area of the head and neck, account for approximately 40% of all cases seen in major pain clinics.

## 5. Conclusion

Pain in any kind is unbearable. When it is oral pain, it is a worse scenario. This condition has to be diagnosed and treated at the earliest. Proper knowledge and clinical skills have to be used in patients. In dental treatment, our goal is to cure the patient of pain. In prosthodontics again same goal of pain including Orofacial aspects of pain also. It has to be first diagnosed and examined carefully. Pain is checked if it is of dental or non dental origin. Then the suitable treatment planning is done. It is important that the patient is also informed and motivated for the treatment. Patients also need encouragement and counselling from dentists and prosthodontists. Depending upon the cause, the Orofacial pain is treated. It requires utmost patience and perseverance from the dentist and prosthodontist as it is time consuming sometime. Nevertheless it is challenging but definitely rewarding.

## 6. Source of Funding

No financial support was received for the work within this manuscript.

## 7. Conflict of Interest

None declared.

## References

1. Black JA, Cummins TR, Plumpton C, Chen YH, Hormuzdiar W, Clare JJ, et al. Upregulation of a silent sodium channel after peripheral, but not central, nerve injury in DRG neurons. *J Neurophysiol.* 1999;82(5):2776–85. doi:10.1152/jn.1999.82.5.2776.
2. Okacle D. A review on oral and dental pain. *J Adv Clin Res Insights.* 2019;6(2):43–7.
3. Goraka PM. Odontogenic pain. *Rad 507 Med Sci.* 2010;34:43–54.
4. Renton T. Dental (odontogenic) pain. *Rev Pain.* 2011;5(1):2–7. doi:10.1177/2049463711100500102.
5. Abbott P. Medical management of dental and oral pain. *Aust Prescr.* 2007;30:77–9. doi:10.18773/austprescr.2007.044.
6. Kaur A, Dhillon N, Singh S, Gambhir RS. Orofacial pain - an update on diagnosis and management. *JMR.* 2017;3(2):93–8.
7. Faguda K. Diagnosis and treatment of abnormal dental pain. *J Dent Anesth Pain Med.* 2016;16(1):1–8.
8. Macauley Y, O'Donnell P, Duncan HF. Dental pain. *BMJ.* 2013;347:f6539. doi:10.1136/bmj.f6539.
9. Okeson JP. Bell's orofacial pains. The clinical management of orofacial pain. Carol stream, IL: Quintessence publishing co, inc; 2005.
10. Okeson JP. The classification of orofacial pains. *Oral Maxillofac Surg Clin North Am.* 2008;20(2):133–44.
11. Niv D. Pain and quality of life. *Pain Pract.* 2001;1(2):150–61. doi:10.1046/j.1533-2500.2001.01016.x.
12. Sally C. Aide. memoires in oral diagnosis: mnemonics and acronyms (the Scully system). *J Investig Clin Dent.* 2012;3(4):262–3. doi:10.1111/jicd.12005.

13. Gupta R, Mohan V, Mahay P, Yadav PK. Orofacial pain - a review. *Dentistry*. 2016;6(3):1–6. doi:10.4172/2161-1122.1000367.
14. Kim K. Non odontogenic toothache caused by acute maxillary sinusitis :a case report. *J Oral Med Pain*. 2016;41(2):80–4.
15. Khatib MS. A review of orofacial pain. *J Adv Clin Res Insights*. 2018;5(6):192–5. doi:10.15713/ins.jcri.242.
16. Young A. OFP overview:getting rid of the riddles. *CDA J*. 2016;44(12):729–35.
17. Van Deun L, de Witte M, Goessens T, Halewyck S, Ketelaer M, Matic M, et al. Facial pain:a comprehensive review and proposal for a pragmatic diagnostic approach. *Eur Neurol*. 2020;83(1):5–16. doi:10.1159/000505727.
18. Abdelnabi AH, Swelem AA. Influence of defective complete dentures renewal on tmd. *Gerodontology*. 2015;32(3):211–21. doi:10.1111/ger.12102.
19. Johansson A, Omar R, Carlsson GE. Bruxism and prosthetic treatment. *J Prosthodont Res*. 2011;55(5):121–36. doi:10.1016/j.jpor.2011.02.004.
20. Agerberg G, Sandstro R. Frequency of occlusal interferences: A clinical study in teenagers and young adults. *J Prosth Dent*. 1988;59(2):212–7.
21. Plesh O. Prosthetic rehabilitation in tmd and Orofacial pain patients. Clinical problem solving. *Dent Clin North Am*. 1992;36(3):581–9.
22. Hagag G, Yoshida K, Miura H. Occlusion, prosthodontic treatment, temporomandibular disorders. *J Med Dent Sci*. 2000;47(1):61–6.
23. Hegarty A, Zakrzewska JM. Differential diagnosis for orofacial pain, including sinusitis, TMD, trigeminal neuralgia. *Dent Update*. 2011;38(6):396–400. doi:10.12968/denu.2011.38.6.396.
24. Zakrzewska JM. Differential diagnosis of facial pain and guidelines for management. *Br J Anaesth*. 2013;111(1):95–104. doi:10.1093/bja/aet125.
25. The scope of TMD/orofacial pain (head and neck pain management) in contemporary dental practice. Dental Practice Act Committee of the American Academy of Orofacial Pain. *J Orofac Pain*. 1997;11(1):78–83.
26. Carrara SV, Cont PCR, Barbosa JS. Prosthodontic planning in patients with tmd/bruxism. *Dent Press J Orthod*. 2010;15(3):114–20.
27. Lipton R, Newman LC. The epidemiology and impact of migraine. *Neurology*. 2003;60(7):S3–8.
28. Roshtamkhani F, Tayarani N, Madani AS. The Relationship between Temporomandibular Disorders (TMDs) and Overall Denture Conditions in Complete Denture Wearers. *JDMT*. 2015;4(2 - Serial Number 2):101–10. doi:10.22038/JDMT.2015.4128.

### Author biography

**Isha Rastogi**, Associate Professor

**Cite this article:** Rastogi I. Assessment of oral, dental and facial pain in patients. *IP Ann Prosthodont Restor Dent* 2022;8(4):216-219.